arXiv:1209.6499v1 [quant-ph] 28 Sep 2012

Simultaneous Estimation of Dimension, States and Measurements: Rigidity Considerations

Cyril Stark[1]

[1]*Institute for Theoretical Physics, ETH Zürich, 8093 Zürich, Switzerland*

The Gram matrix of the experimentally prepared states and the performed measurements determines the associated density matrices and POVM (projective operator valued measure) elements uniquely up to simultaneous rotations of all density matrices and POVM elements. In [Stark, arXiv:1209.5737 (2012)] we showed how the state-measurement Gram matrix can be estimated efficiently, and we have seen that in the generic case involving full-rank density matrices and full-rank POVM elements, the measurement data never suffices to uniquely determine the state-measurement Gram matrix. In this paper, we first provide a sufficient criterion to test, which additional assumptions (such as the projectiveness of some measurements) suffice to guarantee the uniqueness of the state-measurement Gram matrix up to discrete symmetries. Subsequently, inspired by the non-uniqueness of the Gram matrix in situations involving full-rank matrices, we examine more closely scenarios involving projective, non-degenerate measurements and their pure post-measurement states. We find that these experiments can be identified on the sole basis of the collected measurement data. Moreover, the Gram matrix is determined uniquely and can be read off directly from the measurement data.

In a recent paper [1], we introduced a coherent strategy to simultaneously estimate Hilbert space dimension, states and POVMs. More precisely, we showed how one can determine the Gram matrix associated with the involved density matrices and POVM elements. Up to simultaneous rotations of all states and POVM elements in the space of Hermitian matrices, this Gram matrix uniquely determines the family of prepared states and the performed POVMs uniquely. For situations in which knowing the states-measurements Gram matrix is not sufficient, we have developed a heuristic algorithm [2]—taking the Gram matrix as input—to find explicit realizations of the performed states and POVM elements.

In [1], we observed that generically, the measurement data does not uniquely specify the states-measurement Gram matrix—a necessary condition for the uniqueness (up to simultaneous conjugation with unitary or anti-unitary mappings) of the density matrices and POVM elements describing the experiment. The purpose of this paper is the exploration of the set of Gram matrices which are compatible with the experimentally collected measurement data. More precisely, after having introduced the setting in Section I, we describe in Section II a method to test whether specific assumptions (such as the projectiveness of some measurements or the purity of some states) are sufficient to guarantee the uniqueness of the estimated states-measurement Gram matrix G (up to discrete symmetries). Subsequently, in Section III, we describe how the outcome of this test can be predicted with probability one before we have even computed the full states-measurements Gram matrix. This allows for the computation of phase diagrams that are universally applicable. In Section IV we provide some simple examples. In [1] we addressed problems that might arise from missing characterizations of the set of Gram matrices that can be generated via density matrices and POVM elements. In Section V, making use of the uniqueness test, we add a few remarks to this discussion.

The counterexample in [1] showing that generically, the measurement data alone does not uniquely specify the Gram matrix, is not applicable to situations in case of which the family of prepared states as well as the family of performed measurements contain matrices which are rank deficient. This inspired the closer investigation of data tables associated with pure states and projective measurements in Section VI. We show that data tables which are generated by projective, non-degenerate measurements and their associated post-measurement states can be recognized as being generated in this manner. In these cases, the states-measurement Gram matrix can be directly read off from the measurement data. No computations are necessary and the Gram matrix is uniquely determined by the measurement data.

Other works associated with full quantum modeling going beyond state or measurement tomography focused on the falsification of incorrect assumptions about the performed measurements [3], temporal drifts in the devices [4], and consequences of inappropriate quantum modeling [5–7]. On the side of dimension estimation, the works [8–10] investigate linear dimension witnesses to estimate the dimension of the Hilbert space associated with the performed experiment.

I. ESTIMATING EFFECTIVE MODELS VIA CONVEX RELAXATION

In this section, we are introducing the notation and sketch results from [1] about the assumption-free inference of quantum models via state-measurement Gram matrices. To construct quantum models solely based on the measurement data, we are guided by Occam's razor, i.e., we are aiming at finding the simplest quantum model describing the measurement data $(f_{w,vk})_{w,v,k}$. Here, $f_{w,vk}$ denotes the frequency of measuring outcome 'k' in case we have prepared an unknown state ρ_w and measured an unknown POVM $(E_{vk})_k$. More precisely, among all the quantum models which are compatible with the measurement data, we search for those quantum models whose Hilbert space dimension is minimal. We are focusing on the asymptotic regime (i.e., the number of independently repeated measurements goes to infinity), so that according to Born's rule

$$p_{w,vk} = \operatorname{tr}(E_{vk}\rho_w) = f_{w,vk}.$$

Expanding all the unknown states and POVM elements with respect to an orthonormal basis in the space of Hermitian matrices, we can express all states and POVM elements in terms of real (the space of Hermitian matrices forms a real vector space) vectors in $\mathbb{R}^{d^2}$. These vectors can be concatenated into the matrix

$$P = (\vec{\rho}_1 \,|\, \cdots \,|\, \vec{\rho}_W \,|\, \vec{E}_{11} \,|\, \cdots \,|\, \vec{E}_{1K} \,|\, \cdots \,|\, \vec{E}_{V1} \,|\, \cdots \,|\, \vec{E}_{VK}) \tag{1}$$

and

$$G = P^T P. \tag{2}$$

G is the Gram matrix associated with ρ_1, ..., E_{VK}. Here, W and V denote the total number of prepared states and the total number of performed measurements. To keep the notation simple, we assume that each measurement has K outcomes. This assumption leaves the explanations unchanged. We observe that the measurement data appears as off-diagonal block in G:

$$G = \left(\begin{array}{c|c} G_{\mathrm{st}} & \mathcal{D} \\ \hline \mathcal{D}^T & G_{\mathrm{m}} \end{array}\right) \tag{3}$$

where

$$\mathcal{D} = \begin{pmatrix} f_{1,(1,1)} & \cdots & f_{1,(V,K)} \\ \vdots & & \vdots \\ f_{W,(1,1)} & \cdots & f_{W,(V,K)} \end{pmatrix}. \tag{4}$$

Since $\operatorname{rank}(G) = \operatorname{rank}(P)$, finding the minimal quantum model with dimension d is equivalent to finding a Gram matrix G of minimal rank which is compatible (in the sense of (3)) with the measurement data. More precisely, the Gram matrix G associated with the simplest quantum model is a solution of the optimization problem

$$\begin{array}{ll} \text{argmin} & \operatorname{rank} G \\ \text{subject to} & G \in \mathcal{G}_{\mathrm{QM}}, \\ & G \text{ satisfies (3) for given } \mathcal{D}. \end{array} \tag{5}$$

Here, we use $\mathcal{G}_{\mathrm{QM}}$ to denote the set of Gram matrices that can be generated via density matrices and POVM elements. Unfortunately, the rank function is non-convex and consequently, the optimization problem (5) is non-convex. Thus, it suffers from local minima. It has even been shown that rank minimization is NP-hard [11]. To approximately solve problem (5), one can replace the non-convex objective function by its so called 'convex envelope' which is—roughly speaking—the 'closest' convex function. This procedure is called convex relaxation and it maps the original, uncomputable optimization problem to the 'closest' convex optimization problem. In [1], we showed that the trace of positive semidefinite matrices is (up to an irrelevant constant factor), the convex envelope of their rank. Hence, convex relaxation of the uncomputable problem (5) leads to

$$\begin{array}{ll} \text{argmin} & \operatorname{tr} G \\ \text{subject to} & G \geq 0,\ \|G\| \leq M_{\mathrm{QM}}, \\ & G \text{ satisfies (3) for given } \mathcal{D}, \end{array} \tag{6}$$

(the bound M_{QM} plays a minor role). Up to simultaneous Hilbert-Schmidt rotations of all states and POVM elements, G determines the states and the POVM elements uniquely. However, states-measurement configurations computed in this way do not necessarily correspond to positive semidefinite matrices, even though all pairwise inner products satisfy all constraints dictated by G. To find explicit, quantum mechanically valid density matrices and POVM elements, we have developed a heuristic algorithm [2] that takes the Gram matrix G as input and searches for positive semidefinite matrices satisfying all the G-constraints. After having mapped the original rank minimization (5) onto its convex relaxation (6), we have addressed in [1] the question whether or not the Gram matrices (3) are uniquely determined by the measurement data $\mathcal{D}$. A simple counterexample showed that this is not the case if all the involved states and POVM elements are not rank-deficient.

II. FINDING SUFFICIENT CONDITIONS FOR UNIQUENESS

We have seen that data $\mathcal{D}$ generated by full-rank states and measurements does not uniquely specify the states-POVM-configration. Thus, if we are only interested in unique solutions, we need to make additional assumptions about the states-measurement-configuration to guarantee uniqueness. For instance, one could have reasons to assume that some of the measurements are projective or that some of the states are pure. Which assumptions guarantee uniqueness of the states-measurements Gram matrix? Answering this question is the goal of this section. Recall that G is the states-measurement Gram matrix. Assumptions like the projectiveness of measurements can be described by fixing some entries of G. For instance, assume that the measurement "v" is projective. Then POVM elements E_{vk} and E_{vl}, $k \neq l$, are orthogonal and the G-entry corresponding to the inner product between E_{vk} and E_{vl} has to be set to zero. Define Ω to be the set of G-indices that are assumed to be known. Thus, G_Ω represents our knowledge about G.

As mentioned earlier (cf. Eq. (1)), we can rewrite the density matrices and the POVM elements in terms of vectors that themselves build up a matrix P such that $G = P^T P$. When is G uniquely determined by our knowledge G_Ω? We are discussing this question regarding the states and the POVM elements as arbitrary Hermitian matrices, thus ignoring for example the positivity constraint of quantum mechanics. These constraints narrow down the set of possible states and measurement patterns. Consequently, uniqueness without the consideration of the quantum mechanical constraints implies uniqueness under the consideration of the quantum mechanical constraints. We can thus view the column vectors of the states-measurement matrix

$$P = [\vec{p}_1 ... \vec{p}_n] \tag{7}$$

as arbitrary vectors in $\mathbb{R}^R$, with

$$R = \mathrm{rank}(G). \tag{8}$$

In Eq. (8), we implicitly assume that $n \geq R$ because $G = P^T P$ and therefore, $\mathrm{rank}(G) = \mathrm{rank}(P)$. However, demanding that $n \geq R$ is clearly necessary, because otherwise the columns of P could not even span the column vector space $\mathbb{R}^R$ associated with the Hilbert-Schmidt space carrying the density matrices and POVM elements. Moreover, we assume that P has been drawn from a non-singular probability distribution. The following discussion cannot answer the uniqueness question fully but it provides a sufficient condition for the exclusion of smooth deformations of those states-measurement configuration P which are compatible with our knowledge G_Ω. Hence, we are not able to exclude that the set of $\mathcal{D}$-compatible Gram matrices is a set of discrete points. The existence of smooth deformations is equivalent to the existence of an η-dimensional smooth submanifold $N \subseteq \mathbb{R}^{R\times n}$ ($\eta \geq 1$) in an open neighborhood of P, $P \in N$, with the defining property that

$$f^{(k,l)}(Q) = \vec{q}_k^{\,T} \vec{q}_l - G_{kl} = 0 \tag{9}$$

for all $Q = [\vec{q}_1, ..., \vec{q}_n] \in N$ and $(k,l) \in \Omega$. The dimension of N given by

$$\dim(N) = Rn - \mathrm{rank}\left(\vec{\nabla} f^{(k_1,l_1)}(P), ..., \vec{\nabla} f^{(k_{|\Omega|},l_{|\Omega|})}(P)\right) \tag{10}$$

The matrix in the argument on the LHS is called *completion matrix* [12]. We are going to abbreviate it with $C_\Omega(P)$. The following discussion follows remarks of A. Singer and M. Cucuringu in [12] and aims at adapting a discussion conducted in [13] to our setting. Note that N is never zero-dimensional because the left-multiplication of P with elements of $SO(R)$ (i.e., simultaneous rotation of all the columns in P; relative inner products are unchanged) produces new and valid configurations out of P. Let $\tilde{N} \subseteq N$ denote the submanifold generated by these simultaneous rotations of the columns of P. Thus, the existence of non-trivial smooth deformations becomes equivalent to $N \neq \tilde{N}$. Let (φ_N, U_N) and $(\varphi_{\tilde{N}}, U_{\tilde{N}})$ denote the local charts of N and $\tilde{N}$ with $\varphi_N^{-1}(P) \in U_N \subseteq \mathbb{R}^{\dim(N)}$, $\varphi_{\tilde{N}}^{-1}(P) \in U_{\tilde{N}} \subseteq \mathbb{R}^{\dim(\tilde{N})} \subseteq \mathbb{R}^{\dim(N)}$ and $\varphi_N^{-1}(P) = \varphi_{\tilde{N}}^{-1}(P)$, when appropriately embedding $\mathbb{R}^{\dim(\tilde{N})}$ into $\mathbb{R}^{\dim(N)}$. Let $\pi_{\tilde{N}} : \mathbb{R}^{\dim(N)} \to \mathbb{R}^{\dim(N)}$ denote the orthogonal projection onto $\mathrm{span}(U_{\tilde{N}})$. Since U_N and $U_{\tilde{N}}$ are open neighborhoods of $\varphi_N^{-1}(P)$ there exists an $\varepsilon > 0$ such that

$$\pi_{\tilde{N}}(U_N \cap B_\varepsilon(\varphi_N^{-1}(P))) = U_{\tilde{N}} \cap B_\varepsilon(\varphi_N^{-1}(P)).$$

The existence of smooth deformations is equivalent to the existence of points $\vec{x} \in \big(U_N \cap B_\varepsilon(\varphi_N^{-1}(P))\big) \setminus \big(U_{\tilde{N}} \cap B_\varepsilon(\varphi_N^{-1}(P))\big)$. Such vectors $\vec{x}$ exist if and only if there are elements in $U_N \cap B_\varepsilon(\varphi_N^{-1}(P))$ such that their orthogonal projection onto $\ker(\pi_{\tilde{N}}(U_N)$ is non-vanishing. This on the other hand is equivalent to $\dim(U_N) > \dim(U_{\tilde{N}})$ which is equivalent to $\dim(N) > \dim(\tilde{N})$. We conclude that the existence of non-trivial smooth deformations is equivalent to $\dim(N) > \dim(\tilde{N})$. Eq. (10) provides a formula for the dimension of N. In order to check whether or not $\dim(N) > \dim(\tilde{N})$, we are thus left with the task to determine the dimension of $\tilde{N}$.

The dimension of $\tilde{N}$ is equal to the dimension of its tangent space $T_P\tilde{N}$ at P,

$$T_P\tilde{N} = \mathrm{span}\{\dot{Q}(0) \mid Q(t) = \exp(\alpha(\vec{t}) \cdot \vec{I})P\},$$

with $\vec{\alpha}(\cdot) \subset \mathbb{R}^{\dim(SO(R))}$, $\vec{\alpha}(0) = 0$ arbitrary. Here, $\vec{I}$ carries all the generators of $SO(R)$. Rewriting $P = USV^*$,

$$S = \left(\begin{array}{ccc|ccc} s_1 & & & 0 & \cdots & 0 \\ & \ddots & & \vdots & \ddots & \vdots \\ & & s_R & 0 & \cdots & 0 \end{array}\right) \in \mathbb{R}^{R\times n}, \tag{11}$$

$U \in O(R)$, $V \in O(n)$, in terms of its singular value decomposition and using that the product of orthogonal matrices is an orthogonal matrix, we arrive at

$$T_P\tilde{N} = \mathrm{span}\{\dot{W}(0) \mid W(t) = \exp(\beta(\vec{t}) \cdot \vec{I})SV^*\},$$

with $\vec{\beta}(\cdot) \subset \mathbb{R}^{\dim(SO(R))}$, $\vec{\beta}(0) = 0$ arbitrary. Taking the derivative, we get $\dot{W}(0) = (\dot{\vec{\beta}}(0) \cdot \vec{I})P$. The matrix $\dot{\vec{\beta}}(0) \cdot \vec{I}$ is an arbitrary skew-symmetric matrix because $\dot{\vec{\beta}}(0) \in \mathbb{R}^{\dim(SO(R))}$ can be arbitrary. Denote the linear subspace of skew-symmetric $(R \times R)$-matrices by $\mathcal{A} \subset \mathbb{R}^{R\times R}$. We conclude that

$$\dim(\tilde{N}) = \dim\ \mathcal{A}SV^*.$$

The right-action of S stretches $\mathcal{A}$ and embeds it into $\mathbb{R}^{R\times n}$. The action of V^* then simply turns the rows of the resulting matrices according to V. Consequently,

$$\dim(\tilde{N}) = \dim(\mathcal{A}SV^*) = \dim(\mathcal{A}) = \frac{1}{2}R(R-1).$$

We conclude that the *absence* of non-trivial smooth deformations is equivalent to $\dim(N) = \dim(\tilde{N})$ which is itself equivalent to

$$\mathrm{rank}\,(C_\Omega(P)) = Rn - \frac{1}{2}R(R-1). \tag{12}$$

III. A PRIORI CHECK OF SMOOTH FLEXIBILITY

Let $\mathcal{C} := \mathbb{R}^{R\times(W+VK)}$ be the space carrying all states and measurement configurations and set $\kappa := Rn - R(R-1)/2$ (the RHS of Eq. (12)). Following a proof sketch in [12], we are going to show next how Eq. (12) can be tested before we have even computed any states-measurement configuration P that is compatible with the constraints G_Ω. More precisely, we have the following procedure in mind (see Algorithm 1): Fix Ω and draw $Q \in \mathcal{C}$ at random with respect to a non-singular probability measure on $\mathcal{C}$. We claim that then, with probability one, $\operatorname{rank}(C_\Omega(P)) = \kappa$ if and only if $\operatorname{rank}(C_\Omega(Q)) = \kappa$, whenever $P \in \mathcal{C}$ is drawn from an arbitrary non-singular probability measure. Thus, by testing (12) on a randomly sampled Q, we can test the flexibility of P without even knowing P and G_Ω. The knowledge about Ω alone is sufficient.

We start by showing that $\operatorname{rank}(\tilde{P}) = R$ (recall our assumption $n \geq R$) on almost all $\tilde{P} \in \mathcal{C}$. Let $\tilde{P} = USV^*$ be the singular value decomposition of $\tilde{P}$. If we restrict ourselves to matrices $\tilde{P}$ with the property that the number of non-zero singular values appearing in S is smaller than R then we are loosing degrees of freedom. Thus, we are constrained to a lower-dimensional submanifold embedded into $\mathcal{C}$. Consequently, almost all elements in $\mathcal{C}$ have full rank. So, both P and Q from before have full rank and we know the value of κ a priori with probability one.

Next, we are proving the asserted relation between the flexibility of P and the flexibility of Q by showing that the answer to the question whether or not condition (12) is fulfilled is the same for almost all matrices in $\mathcal{C}$. Let

$$\mathcal{N} := \{Q \in \mathcal{C} \mid \operatorname{rank}(C_\Omega(Q)) = \kappa\}$$

Proving the claim is equivalent to showing that $\mathcal{N}$ is either equal to $\mathcal{C}$ or it is a zero-set in $\mathcal{C}$. Define

$$\omega_\kappa(Q) := \sum_{\substack{A \subseteq C_\Omega(Q)\\ A \in \mathbb{C}^{\kappa\times\kappa}}} (\det A)^2$$

($A \subseteq C_\Omega(Q)$ means that A is a submatrix of $C_\Omega(Q)$). We arrive at the auxiliary claim that

$$\mathcal{N} = \{Q \in \mathcal{C} \mid \omega_\kappa(Q) \neq 0\} \tag{13}$$

"$\subseteq$": Since $\operatorname{rank}(C_\Omega(Q)) = \kappa$ there exists a non-zero minor of size κ.
"$\supseteq$": By assumption there exist at least one non-zero $C_\Omega(Q)$-minor of size κ. Thus, $\operatorname{rank}(C_\Omega(Q)) \geq \kappa$. It remains to show that $\operatorname{rank}(C_\Omega(Q)) > \kappa$ is impossible. This is indeed the case because $\operatorname{rank}(C_\Omega(Q)) > \kappa$ would imply that

$$\begin{aligned} Rn - \kappa &= \dim(\tilde{N}) \\ &\leq \dim(N) = Rn - \operatorname{rank}(C_\Omega(Q)) < Rn - \kappa. \end{aligned} \tag{14}$$

This is the desired contradiction.

From Eq. (13) we conclude that $\mathcal{C}\backslash\mathcal{N}$ is the set of roots of a polynomial in the entries of $Q \in \mathcal{C}$. The polynomial is determined by its values on a finite set of points. Thus, whenever $\mathcal{C}\backslash\mathcal{N}$ is not a zero-set, $\omega_\kappa(Q)$ agrees with the zero function on a set with accumulation point, and therefore, $\omega_\kappa(Q)$ is equal to the zero function. But then, $\mathcal{C}\backslash\mathcal{N} = \mathcal{C}$. Consequently, either $\mathcal{C}\backslash\mathcal{N}$ is a zero set, or $\mathcal{C}\backslash\mathcal{N} = \mathcal{C}$. Thus, we have shown that by testing (12) on a randomly sampled Q, we can test the flexibility of P without knowing neither P nor G_Ω; knowing Ω is sufficient. We arrive at Algorithm 1 for the a priori test of flexibility. Recall that our analysis allowed arbitrary matrices $P \in \mathbb{R}^{R\times n}$ and did not take into account quantum mechanical constraints such as the positive-semidefiniteness of density matrices and POVM elements. Such constraints can constrain the flexibility of P. Therefore, Algorithm 1 does not permit any conclusion about flexibility whenever the condition (12) is not fulfilled. Algorithm 1 provides a sufficient and not a necessary criterion for the exclusion of smooth deformability.

Algorithm 1 (Cucuringu, Singer [12])

```
Require: $\Omega$, $R = \operatorname{rank}(G)$, $n = W + VK$
Draw $Q$ at random from a non-singular distribution.
Compute the completion matrix $C_\Omega(Q)$.
If $\operatorname{rank}(C_\Omega(Q)) = Rn - \frac{1}{2}R(R-1)$ then, with probability
    one, almost all $P \in \mathcal{C}$ are not deformable.
```

IV. RIGIDITY PHASE DIAGRAMS

For each tupel (W, V, K, Ω, R), Algorithm 1 determines the property 'locally rigid' or 'locally flexible'. The results Algorithm 1 produces when varying W and V, can be merged to form a rigidity phase diagram (see Figure 1). In the following, when drawing rigidity diagrams, we are each time setting R (equal to d^2) to a specific value and start to vary the number of states W and the number of measurements V. Points in these diagrams that are incompatible with the constraint $n \geq R$, $R = d^2$, are automatically assigned to the deformable phase; n denotes the number of columns of P from Eq. (7).

We have already observed that $\mathcal{D}$ by itself does not lead to rigidity. To achieve rigid states-measurement configurations we have to impose more constraints on G. Taking seriously the outstanding role of energy measurements in experiments, we might promote these measurements to be projective and non-degenerate. But we did not manage to find a rigid phase given this assumption (e.g., for $\dim\mathcal{H} = 2$, the point $(W, V, K) = (100, 100, 2)$ is still in the non-rigid phase). Each projective, non-degenerate measurement provides—via the associated post-measurement states—a set of $\dim\mathcal{H}$ orthonormal pure states. But even when postulating the energy post-measurement states to be pure and orthogonal, we failed



FIG. 1: Rigidity phase diagram in case of $\dim \mathcal{H} = 2$ and pure states.



FIG. 2: Rigidity phase diagram in case of $\dim \mathcal{H} = 2$ and projective measurements.

finding a rigid phase. As a further tightening, we try out the assumption that all involved states are pure (now letting loose of the projective energy measurement). Figure 1 shows the rigidity phase diagram in case of two dimensional systems.

We conclude that there exists a rigid phase in case the experimentalist knows the Hilbert-Schmidt norms of the measured states. Figure 2 shows the rigidity phase diagram in case one correctly postulates that all the measurements are projective and non-degenerate. This condition is even stronger than the condition that all states are pure because the projectiveness of non-degenerate measurements implies orthogonality of the POVM elements. Thus, in addition to Hilbert-Schmidt norms of the POVM elements (diagonal elements in G), we also know those off-diagonal elements of G that describe the inner product between different elements of the same POVM.



FIG. 3: Rigidity phase diagram in case of $\dim \mathcal{H} = 2$, pure states and projective, non-degenerate measurements.



FIG. 4: Rigidity phase diagram in case of $\dim \mathcal{H} = 3$, pure states and projective, non-degenerate measurements.

Figures 3, 4 and 5 display rigidity phase diagrams when postulating the combination of the preceding two scenarios, i.e., pure states and projective, non-degnerate measurements.

V. ISSUES IN CONNECTION WITH $\mathcal{G}_{\mathrm{QM}} \neq S^{+} \cap B_{\|\cdot\|_{\mathrm{op}} \leq M_{\mathrm{QM}}}$

To simultaneously estimate dimension, states and measurements we first estimate the state-measurement Gram matrix via the optimization problem (6). Here, we have to include the entire knowledge G_Ω as constraints. Afterwards, we compute the rank of G and check via Algorithm 1 whether or not G lies within the rigid phase. However, we are not able to check whether or not $G \in \mathcal{G}_{\mathrm{QM}}$ (cf. Eq. (5)). If we believe that the data was gener-



FIG. 5: Rigidity phase diagram in case of $\dim \mathcal{H} = 4$, pure states and projective, non-degenerate measurements.

ated by a quantum model with Hilbert space dimension equal to $\sqrt{\mathrm{rank}(G)}$ then $G \in \mathcal{G}_{\mathrm{QM}}$ whenever G is in the rigid phase because there is simply no other constraint-compatible Gram matrix with that rank (when ignoring hypothetical discrete symmetries). However, as long as we are missing a precise characterization of $\mathcal{G}_{\mathrm{QM}}$ or the dimension of the underlying Hilbert space, we do not know how to guarantee easily that the simplest quantum Gram matrix compatible with (26) does not have higher rank then the computed matrix G. This possibility can be excluded when we are able to find explicit density matrices and POVM elements reproducing G. If we manage to find explicit density matrices and POVM elements, then we are sure that G corresponds to a quantum model, the rigidity considerations apply and we can be sure to have found the unique simplest quantum Gram matrix explaining the data. A heuristic method (taking G as input) to compute explicit density matrices and POVM elements on the basis of a Gram matrix will be presented in [2].

VI. CERTIFIED UNIQUENESS IN CASE OF PROJECTIVE, NON-DEGENERATE MEASUREMENTS

Recall the 2×2 block matrix structure

$$G = \begin{pmatrix} G_{\mathrm{st}} & \mathcal{D} \\ \mathcal{D}^T & G_{\mathrm{m}} \end{pmatrix} \tag{15}$$

of the states-measurements Gram matrix G (cf. Eq.(3)). Assume that the measurement 't' described by the POVM $(E_{tk})_{k=1}^d$ is projective and non-degenerate. Thus, it is described by d orthogonal rank-one projectors. The post-measurement states $(\rho_{tw})_{w=1}^d$ associated with the measurement $(E_{tk})_{k=1}^d$ are d orthonormal pure states. Let $\mathcal{D}_\square$ denote the submatrix of $\mathcal{D}$ associated with the post-measurement states $(\rho_{tw})_{w=1}^d$ and the projective measurement $(E_{tk})_{k=1}^d$. We conclude that

$$\mathcal{D}_\square = \mathbb{I}_d. \tag{16}$$

Eq. (16) forms a necessary condition for $(E_{tk})_{k=1}^d$ to be projective and non-degenerate. Let $G_\square$ denote the $(2d \times 2d)$-submatrix of G associated with $\mathcal{D}_\square$, so that

$$G_\square = \begin{pmatrix} (G_{\mathrm{st}})_\square & \mathcal{D}_\square \\ \mathcal{D}_\square^T & (G_{\mathrm{m}})_\square \end{pmatrix} \geq 0. \tag{17}$$

Next, we are going to show that Eq. (16) forms a necessary and sufficient condition for $(E_{tk})_{k=1}^d$ to be projective and non-degenerate, and for $(\rho_{tw})_{w=1}^d$ to be the associated pure post-measurement states. By Cauchy-Schwarz inequality,

$$\|\rho_j\|_2^2 \, \|E_{tj}\|_2^2 \geq |\mathrm{tr}(E_{tj}\rho_j)| = 1. \tag{18}$$

Among all states, the pure states have maximal Hilbert-Schmidt norm, so that

$$\|\rho_j\|_2^2 \leq 1. \tag{19}$$

Eq. (18) implies that

$$\|E_{tj}\|_2^2 \geq 1. \tag{20}$$

On the other hand, we know that $(E_{tk})_{k=1}^d$ is a POVM. Therefore, $\sum_j E_{tj} = \mathbb{I}$ and $E_j \geq 0$. It follows that

$$\begin{aligned} d = \|\mathbb{I}\|_2^2 = \left\| \sum_{j=1}^d E_{tj} \right\|_2^2 &= \sum_{j,\tilde{j}=1}^d \mathrm{tr}(E_{tj}E_{t\tilde{j}}) \\ &= \sum_{j=1}^d \|E_{tj}\|_2^2 + \sum_{j \neq \tilde{j}} \mathrm{tr}(E_{tj}E_{t\tilde{j}}) \end{aligned} \tag{21}$$

Recall that $\mathrm{tr}(MN) \geq 0$ for all $M, N \geq 0$. Using this and Eq. (20) in Eq. (21), we conclude that

$$\begin{aligned} \|E_{tj}\|_2^2 &= 1, \\ \mathrm{tr}(E_{tj}E_{t\tilde{j}}) &= 0, \text{ for all } j \neq \tilde{j}, \end{aligned} \tag{22}$$

and therefore,

$$((G_{\mathrm{m}})_\square)_{j\tilde{j}} = \delta_{j\tilde{j}} \tag{23}$$

for all $j, \tilde{j}$. This in turn implies (cf. Eq. (18)

$$\|\rho_{tj}\|_2^2 \geq 1 \Rightarrow \|\rho_{tj}\|_2^2 = 1, \tag{24}$$

and therefore,

$$((G_{\mathrm{st}})_\square)_{jj} = 1 \tag{25}$$

for all $j \in \{1, ..., d\}$. Consequently, Eq. (16) forms a necessary and sufficient condition for $(E_{tk})_{k=1}^d$ to be projective and non-degenerate.

Assume that all the considered measurements satisfy Eq. (16). Let V and d denote the number of measurements and the Hilbert space dimension respectively. Then, our knowledge of $G \in \mathbb{R}^{2Vd\times 2Vd}$ takes the form

$$G = \left(\begin{array}{ccc|ccc} 1 & * & * & & & \\ * & \ddots & * & & \mathcal{D} & \\ * & * & 1 & & & \\ \hline & & & \mathbb{I}_d & * & * \\ & \mathcal{D}^T & & * & \ddots & * \\ & & & * & * & \mathbb{I}_d . \end{array}\right). \qquad (26)$$

Zooming back into the specific measurement 't' from before using Eq. (16), we arrive at

$$G_{\square} = \left(\begin{array}{ccc|c} 1 & * & * & \\ * & \ddots & * & \mathbb{I}_d \\ * & * & 1 & \\ \hline & \mathbb{I}_d & & \mathbb{I}_d \end{array}\right). \qquad (27)$$

It follows that

$$1 = \mathrm{tr}(\bar{\rho}_{tj}^T E_{tj}) = \|\rho_{tj}\|_2 \, \|E_{tj}\|_2 = 1 \qquad (28)$$

However, equality in Cauchy-Schwarz can only happen if the involved vectors are linearly dependent. Therefore there exists a scalar κ, such that $\rho_{tj} = \kappa E_{tj}$. Hermiticity of ρ_{tj} and E_{tj} implies that κ is real. Taking the norm on both sides of the equation $\rho_{tj} = \kappa E_{tj}$, we get $|\kappa| = 1$. Hence, $\kappa = 1$ because both ρ_{tj} and E_{tj} are positive semidefinite. We conclude that

$$\rho_{tj} = E_{tj} \qquad (29)$$

for all t, j. Therefore,

$$\mathrm{tr}(\rho_{tj} E_{si}) = \mathrm{tr}(\rho_{tj}\rho_{si}) = \mathrm{tr}(E_{tj}E_{si}) \qquad (30)$$

implying that

$$G_{\mathrm{st}} = \mathcal{D} = G_{\mathrm{m}}. \qquad (31)$$

Consequently,

$$G = \begin{pmatrix} \mathcal{D} & \mathcal{D} \\ \mathcal{D} & \mathcal{D} \end{pmatrix}, \qquad (32)$$

showing that $\mathcal{D}$ is the simultaneous Gram matrix of the states and the POVM elements. We conclude that, once Eq. (16) is fulfilled for every measurement, then the data $\mathcal{D}$ uniquely determines the associated Gram matrix via Eq. (32).

VII. CONCLUSIONS

The freedom of choosing the states-measurements Gram matrix G can (up to discrete symmetries) be regarded as flexibility of the matrices generating the Gram matrix G (under the constraints imposed by our knowledge of G). Fixing the rank of G and the index set Ω marking the known G-entries, we showed, using insights from rigidity theory, how we can test a Gram matrix G for its flexibility before we even know the entries of G explicitly. The specification of the index set Ω suffices. Determining the rigidity of G for different numbers of states, measurements and outcomes, we arrived at rigidity phase diagrams. To apply this method, we had to allow for all G being positive semidefinite instead of G being elements of the set $\mathcal{G}_{\mathrm{QM}}$ (the set of Gram matrices which can be generated quantum mechanically). Since all elements in $\mathcal{G}_{\mathrm{QM}}$ are automatically positive semidefinite, being in the rigid phase is a sufficient but not a necessary condition for rigidity.

The counterexample in [1] which showed that $\mathcal{D}$ does not uniquely determine G applies not to experiments involving rank-deficient states and POVM elements. This lead us to study more closely data tables $\mathcal{D}$ associated with projective, non-degenerate measurements, and we observed that these data tables can be recognized as being generated by projective, non-degenerated measurements. Moreover, G can be directly read off from the measurement data. Consequently, the measurement data determines G uniquely and no computations are needed to determine G.

Acknowledgements

I want to thank Johan Åberg for his continuous support and patient explanations. I count myself very lucky having had the opportunity for discussions with Matthias Baur, Matthias Christandl, Dejan Dukaric, Frédéric Dupuis, Philippe Faist, David Gross, Patrick Pletscher, Renato Renner, Lars Steffen, Lídia del Rio, Sharon Wulff, and Mário Ziman. I acknowledge support from the Swiss National Science Foundation through the National Centre of Competence in Research "Quantum Science and Technology".

[1] C. Stark. Simultaneous estimation of dimension, states and measurements: Gram estimations. *arXiv:1205.2300*, 2012.

[2] C. Stark. Simultaneous estimation of dimension, states and measurements: Computation of representative density matrices and POVMs. *In Preparation.*

[3] T. Moroder, M. Kleinmann, P. Schindler, T. Monz, O. Gühne, and R. Blatt. Detection of systematic errors in quantum experiments. *arXiv:1204.3644*, 2012.

[4] S.T. Flammia, D. Gross, L. Yi-Kai, S. Becker, and J. Eisert. Quantum tomography via compressed sensing: Error bounds, sample complexity, and efficient estimators. *arXiv:1205.2300*, 2012.

[5] I. Gerhardt, Q. Liu, A. Lamas-Linares, J. Skaar, V. Scarani, V. Makarov, and C. Kurtsiefer. Experimentally faking the violation of Bell's inequalities. *Phys. Rev. Lett.*, 107:170404, 2011.

[6] A. Acín, N. Gisin, and L. Masanes. From Bell's theorem to secure quantum key distribution. *Phys. Rev. Lett.*, 97:120405, 2006.

[7] D. Rosset, R Ferretti-Schöbitz, J.D Bancal, N. Gisin, and Y.C. Liang. Imperfect measurements settings: implications on quantum state tomography and entanglement witnesses. *arXiv:1203.0911*, 2012.

[8] R. Gallego, N. Brunner, C. Hadley, and A. Acín. Device-independent tests of classical and quantum dimensions. *Phys. Rev. Lett.*, 105:230501, 2010.

[9] M. Hendrych, R. Gallego, M. Micuda, N. Brunner, A. Acín, and J.P. Torres. Experimental estimation of the dimension of classical and quantum systems. *Nature Phys.*, 8:588–591, 2012.

[10] M. Dall'Arno, E. Passaro, R. Gallergo, and A. Acín. Robustness of device independent dimension witnesses. *arXiv:1207.2574*, 2012.

[11] B. Recht, M. Fazel, and P.A. Parillo. Guaranteed minimum rank solutions to linear matrix equations via nuclear norm minimization. *SIAM Review*, 52(53):471, 2010.

[12] A. Singer and M. Cucuringu. Uniqueness of low-rank matrix completion by rigidity theory. *SIAM J. Matrix Anal. Appl.*, 31:1621–1641, 2010.

[13] H. Gluck. Almost all simply connected closed surfaces are rigid. *Geometric Topology, Lect. Notes Math.*, 438:225–239, 1975.